J Biochem. 2017 Jan;161(1):87-97. doi: 10.1093/jb/mvw057. Epub 2016 Oct 13.

# A novel L-fucose-binding lectin from Fenneropenaeus indicus induced cytotoxicity in breast cancer cells

Biji Chatterjee [1], Krishna Ghosh [1], Nitin Yadav [1], Santosh R Kanade [2]

Affiliations

PMID: 27742744 DOI: 10.1093/jb/mvw057

## Abstract

Lectins are omnipresent in almost all life forms, being the proteins which specifically bind to carbohydrate moieties on the cell surface; they have been explored for their anti-tumour activities. In this study, we purified a fucose specific-lectin (IFL) from Fenneropenaeus indicus haemolymph using fucose-affinity column and characterized for its haemagglutination activity, carbohydrate specificity, dependency on cations and cytotoxicity against cancer cells. The lectin showed non-specificity against human erythrocytes. It was a $Ca^{2+}$-dependent lectin which remained stable over wide pH and temperature ranges. The lectin showed effective dose dependent cytotoxicity against different human cancer cell lines and induced apoptosis in MCF-7 cells as evidenced by DNA ladder assay and PARP cleavage in a dose dependent manner. Moreover, an increased p21 level corresponding to cyclin D downregulation in response to IFL treatment was observed which might work as probable factors to inhibit cell growth and induce apoptosis of MCF-7 cells. Therefore, we report a novel lectin from the prawn haemolymph with high specificity for L-fucose and antiproliferative towards human cancer cells. However, further establishment of the modus operandi of this lectin is required to enable its biotechnological applications.

**Keywords:** Fenneropenaeus indicus; PARP cleavage; apoptosis; fucose-specific Lectin; p21.



## Related information

MedGen

PubChem Compound (MeSH Keyword)

## LinkOut - more resources

**Full Text Sources**

Ovid Technologies, Inc.
Silverchair Information Systems

**Medical**

Genetic Alliance
MedlinePlus Health Information

**Research Materials**

NCI CPTC Antibody Characterization Program