



## Research paper

# Cd(II) and Ni(II) complexes from aroyl hydrazones: Unravelling the intermolecular interactions and electronic, crystal structures through experimental and theoretical studies



S.S. Sreejith<sup>a</sup>, Anju Nair<sup>a</sup>, Victoria A. Smolenski<sup>b</sup>, Jerry P. Jasinski<sup>b</sup>, M.R. Prathapachandra Kurup<sup>a,c,\*</sup>

<sup>a</sup> Department of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, Kerala, India

<sup>b</sup> Department of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA

<sup>c</sup> Department of Chemistry, School of Physical Sciences, Central University of Kerala, Riverside Transit Campus, Neeleshwar 671 314, Kerala, India

## ARTICLE INFO

## Article history:

Received 31 May 2017

Received in revised form 11 September 2017

Accepted 12 September 2017

Available online 21 September 2017

## Keywords:

Crystal structures

Aroyl hydrazones

Hirshfeld surface analysis

ESP

DFT studies

Dipole moment

## ABSTRACT

Three complexes of Cd(II) **1**, **2** and Ni(II), **3** were prepared from two different aroyl hydrozone ligands (HL<sup>1</sup> (C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>4</sub>) & HL<sup>2</sup> (C<sub>20</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>)) using their respective metal bromides. Crystal structures of all the three complexes have been established and the anionic/neutral nature of the coordinating ligand is mainly dictated by the nature of the metal center used. Quantification of intermolecular interactions is done using Hirshfeld surface and 2D finger print analysis. The Electrostatic Potential (ESP) analysis reveals that **2** has a collective concentration of negative electrostatic potentials to one side of the molecule giving a perfect separation of charges. DFT/semiempirical calculations revealed the same, giving a very high dipole moment for the optimized geometry of **2**. Analysis of frontier orbitals is done through DFT calculations at the B3LYP/def2-TZVP level of theory and atomic charge distributions are predicted using AIM method.

© 2017 Elsevier B.V. All rights reserved.

## 1. Introduction

Schiff bases, particularly hydrazones have been active in the synthetic scenario for a long time thanks to the versatility and tunability of coordination modes that they own [1–3]. The degree of metamorphism achieved by these class of ligands and their chelating capability is unmatched by any other. This qualifies them as privileged ligands and aroyl hydrazones are no exception. These ligands and their metal complexes exhibit gripping magnetic [4,5], optical [6,7], catalytic [8,9] and pharmacological applications [10,11]. Out of these myriad of applications, the one that stands out the most is the antibacterial activity [12–14] that they exhibit.

Quantification of intermolecular interactions present in a crystal system has further opened the realm of understanding these interactions. One important tool in this regard is the calculation of a molecular Hirshfeld surface. It is no wonder that the number of publications incorporating this relatively young aspect is increasing daily [15]. A Hirshfeld surface is a single 3D surface that sums up the crystal packing and this 3D surface can further be reduced to a 2D finger plot which is unique for a molecule, thus

deciphering the interactions present in the system [16]. The basic ideas of a Hirshfeld surface and its fingerprint plots are well defined by Spackman et al. [17].

On the other hand, visualization of electrostatic potential surfaces [18–20] can rationalize the intermolecular interactions in the system and the degree of reactivity of the molecules [21]. Apart from the easy visualization of the electrophilic and nucleophilic regions within a chemical species, this mapping will assist in the prediction of reactivity and noncovalent interaction which helps in great detail in dissecting the intermolecular interactions [22,23].

In continuation of our work on aroyl hydrazones and their complexes [24], herein, we report synthesis, characterization and structural study of three new Cd(II) and Ni(II) complexes from two different aroyl hydrazone ligands, HL<sup>1</sup> (C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>4</sub>) and HL<sup>2</sup> (C<sub>20</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>). A Ni(II) metal center, [[Ni(L<sup>2</sup>)<sub>2</sub>·H<sub>2</sub>O], **3**; L = C<sub>20</sub>H<sub>16</sub>N<sub>3</sub>O<sub>2</sub>}, was used in this study since Ni(II) complexes can show interesting biological and electrochemical properties [25]. On the other hand, a Cd(II) metal center, {[Cd(HL<sup>1</sup>)<sub>2</sub>Br<sub>2</sub>], **1** and [Cd(HL<sup>2</sup>)<sub>2</sub>Br<sub>2</sub>], **2**}, is reported to enhance luminescence intensity of the ligand system which can be used for optoelectronic applications [26]. The work described here indicates the versatility in the behavior of the aforementioned ligand systems as in the three complexes synthesized; the ligands exhibit all the possible permutation and combinations of coordination/electronic properties that are

\* Corresponding author at: Department of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, Kerala, India.

E-mail addresses: [mrpcusat@gmail.com](mailto:mrpcusat@gmail.com), [mrp\\_k@yahoo.com](mailto:mrp_k@yahoo.com) (M.R. Prathapachandra Kurup).