

Comparative Secretome Prediction and Analysis of Two *Phytophthora* spp.

Kumar Arvind^{1*}, Rahul C.U.², Merin Babu³, Tony Grace¹ and Rajesh M.K.²

1. Central University of Kerala, Kasaragod, Kerala, INDIA

2. ICAR-Central Plantation Crops Research Institute, Kasaragod, Kerala, INDIA

3. ICAR- Central Plantation Crops Research Institute (RS), Kayamkulam, Kerala, INDIA

*arvind90.psc@gmail.com

Abstract

Phytophthora spp. are widely distributed pathogens causing some of the most devastating diseases in plants. They accomplish parasitic colonization in hosts by virtue of an array of signaling proteins. Secretome analysis is one of the methods to attain molecular insights into species pathogenicity. In this study, we have analyzed the proteome of two species viz., *P. ramorum* and *P. sojae* and predicted extracellular secretory proteins particularly involved in host-pathogen interactions. The potential *Phytophthora* spp. secretome comprising of both the classical and non-classical secretory proteins was predicted with the aid of a stringent computational pipeline.

Out of the 15,743 *P. ramorum* and 19,027 *P. sojae* proteins which were computationally analyzed in the current study, 1396 (8.86%) and 1666 (8.75%) proteins were categorized as classical secretory proteins (CSPs) while a total of 71 (0.45%) and 96 (0.50%) proteins were categorized as non-classical secretory proteins (NCSPs) in *P. ramorum* and *P. sojae* respectively. In addition, 235 and 399 effector proteins were also predicted from *P. ramorum* and *P. sojae* respectively. The functional annotation of the effector proteins revealed the occurrence of SSPs (small specific proteins), virulence and avirulence factors which could prove to be future target to control the pathogenicity and to decipher its role in host-parasite interactions.

Keywords: *Phytophthora* spp., CSPs, NCSPs, SSPs.

Introduction

The genus *Phytophthora* is known worldwide for its pathogenicity on a wide range of plants. This eukaryotic plant pathogenic oomycete causes diseases not only on economically important agricultural crops, but also on ecologically valuable tree species of natural forests¹. Pathogenesis is a dynamic and intricate process modulated by a plethora of extracellular signals secreted by both host plants and pathogen. Pathogens have been known to invade plants by virtue of various classical and non-classical secretory proteins (CSPs and NCSPs)².

The oomycetes accomplish parasitic colonization of plants by reprogramming the defense circuitry of host cells through

an array of disease effector proteins³. Studies on molecular mechanisms underlying *Phytophthora*-host interactions have always elicited an interest in researchers as they could aid in the development of new plant protection strategies. The whole genome sequencing of certain *Phytophthora* spp., along with the development of high-throughput bioinformatics tools, have revolutionized the field of molecular plant- *Phytophthora* interactions. The proteins involved in the host- *Phytophthora* interactions might also play a role in ascertaining the host range⁴.

P. ramorum and *P. sojae* are the two species whose genomes have been sequenced. One of the interesting, but contrasting features of the two species is that *P. sojae* is host-specific whereas *P. ramorum*, has a broad host range. *P. ramorum* incites disease predominantly in forest tree species and poses a major threat to the natural ecological balance. The sudden oak death disease caused by *P. ramorum* affects the keystone species and is causing devastating damage to the ecosystems of Western America^{5,6}. The pathogen has a wide host range affecting plant species from Aceraceae, Anacardiaceae, Betulaceae, Caprifoliaceae, Cupressaceae, Ericaceae, Fagaceae, Hippocastanaceae, Lauraceae, Pinaceae, Rhamnaceae and Rosaceae^{7,8}.

Diseases caused by *P. ramorum* have been reported from USA, Canada, Germany, the Netherlands, Belgium, Denmark, France, Slovenia, Spain, Sweden, England and the Republic of Ireland⁹. *P. sojae* is the causal agent of root rot and stem rot of soybean. The pathogen has a limited host range and zoospores of *P. sojae* are reported to be chemotactically attracted to the isoflavones - daidzein and genistein exuded by soybean roots¹⁰. In this study, we have carried out comparative secretome prediction in these two *Phytophthora* spp. and detailed analyses to investigate effector proteins which could be putatively involved in host-pathogen interactions.

Material and Methods

Secretome prediction: We have used complete predicted proteomes of two *Phytophthora* spp. viz. *P. ramorum* and *P. sojae* which are available on the fungal genomics database developed by the Joint Genome Institute¹¹. The proteomes of these two species were analyzed and compared with the aid of a stringent computational pipeline developed specifically for this study, the schema of which is shown in figure 1. The presence of secretion signals in the sequences was investigated by the virtue of SignalP v.2¹². Further, transmembrane helices were predicted with TMHMM^{13,14}.

The non-classical secretory proteins which are known to lack signal peptides at their N-terminal regions and are called 'leaderless' sequences, were predicted using SecretomeP¹⁵. TargetP¹⁶ was employed for investigating the subcellular localization of the proteins. Non-plant option was chosen for the organism group and the default parameters were set for the tool. The sequences predicted as 'S' (secretory signal peptides) were selected for the further analysis. The total sequences predicted as 'S' were further analyzed to look for the presence of GPI-anchored proteins by the virtues of GPIsom¹⁷. Predicted proteins shorter than 40 amino-acids were excluded from the analysis.

Effector proteins and effector proteins cluster prediction: The total predicted secretory proteins (including classical and non-classical secretory proteins) obtained through the study were further analyzed by the virtue of EffectorP¹⁸ for the prediction of effector proteins. Further, intra-species BLAST search was performed using the standalone BLAST server¹⁹. The predicted effector proteins obtained were annotated using Blast2GO²⁰. Orthologous effector proteins clusters between *P. ramorum* and *P. sojae* were identified using Inparanoid²¹ (standalone version 4.1) while taking into account the certain parameters viz. score cutoff 40 bits; sequence overlap cutoff 0.5; group merging cutoff 0.5 and scoring matrix BLOSUM62.

Results and Discussion

In this study, initially, the presence of extracellular secretory signals was predicted using a combination of SignalP,

TMHMM and TargetP²². Out of the 15,743 proteins analyzed in the *P. ramorum* and 19,027 analyzed in *P. sojae*, 1396 and 1666 proteins were categorized as CSPs (Classical Secretory Proteins). Normally, eukaryotic secretory proteins have been known to follow classical ER-Golgi pathway, involving a characteristic feature of having a short N-terminal signal peptides²³. However, several secretory proteins that lack the signal peptides are found to be exported via a non-classical secretion pathway. Therefore, in order to look for the presence of non-classical secretory proteins, SecretomeP was employed. The results obtained were subjected to GPIsom analysis to investigate the presence of GPI anchored proteins. We have predicted 71 and 96 proteins respectively from *P. ramorum* and *P. sojae* as NCSPs (Non-Classical Secretory Proteins).

Further, total predicted secretory proteins (CSPs and NCSPs) were taken into account for the prediction of effector proteins. A total of 235 and 399 effector proteins were predicted as candidate effectors in *P. ramorum* and *P. sojae* respectively with effector probability >0.5 by virtue of EffectorP as represented in figure 2. Annotation of effector proteins reveals the occurrence of 19 effectors to act as avirulence factors as mentioned in table 1. In this study, a total of 268 and 353 GPI-anchored proteins (GPI-APs) have been reported and predicted to be found on the external leaflet of the membrane in the respective species. An earlier study undertaken in *F. graminearum* suggests that GPI-APs possess a crucial role in growth, development and virulence as well as have the potential to act as diagnostic markers²⁴.




Figure 1: Computational pipeline used for the prediction of secretory proteins (i.e. classical and non-classical proteins).

We have compared the effector proteins predicted in both the species. In order to investigate sequence homology between *Phytophthora* spp., BlastP was performed with e-value set as equal to less than 1 using standalone NCBI-BLAST. The annotation results revealed that a majority of the effector proteins were found to function as virulence factors and toxins to facilitate infection and trigger defense responses based on recognition of avirulence factors and elicitors²⁵. In this study, various elicitors as mentioned in table 2 which could be categorized as pathogen associated molecular patterns (PAMPS) for both the species were investigated.

Elicitins are species-specific avirulence factors which belong to a family of small proteins reported to be secreted by *Phytophthora* and *Pythium* sp. and are classified as oomycete PAMPs having a characteristic molecular feature of sterol-binding^{26,27}. Previously, three elicitors specific to *P. infestans* have been reported, which are known to constitute a powerful strategy for crop protection by inducing defense response in plants²⁸. Plants recognize the PAMPs by the virtue of plant receptors called pattern recognition receptors (PRRs) and get induced to distal systemic resistance^{25,27}.




Figure 2: Graph showing comparative relation between proteomes, predicted secretory proteins (CSPs & NCSPs) and effector proteins in *P. ramorum* and *P. sojae*.




Figure 3: Representation of orthologous and paralogous protein clusters in the two species.

Table 1
Annotation results showing the occurrence of avirulence factors with Seq ID, functional description and the corresponding E-value in (1) *P. ramorum* and (2) *P. sojae*.

S.N.	<i>Phytophthora ramorum</i> ID	Description	E-value
1.	jgi Phyra1_1 79329	Avirulence 1b	4.5E-9
2.	jgi Phyra1_1 81610	Avirulence 1b	9.9E-36
3.	jgi Phyra1_1 74075	Avr1b-1 avirulence	2.1E-22
4.	jgi Phyra1_1 80524	Avr1b-1 avirulence	4.9E-43
5.	jgi Phyra1_1 80530	Avr1b-1 avirulence	2.5E-27
6.	jgi Phyra1_1 80531	Avr1b-1 avirulence	4.7E-25
7.	jgi Phyra1_1 80533	Avr1b-1 avirulence	4.5E-28

(1)

S.N.	<i>Phytophthora sojae</i> ID	Description	E-value
1.	jgi Physo1_1 135585	Avirulence 1b	1.9E-82
2.	jgi Physo1_1 135621	Avr1b-1 avirulence	2.3E-48
3.	jgi Physo1_1 136046	Avirulence 1b	4.6E-87
4.	jgi Physo1_1 136207	Avr1b-1 avirulence	1.2E-80
5.	jgi Physo1_1 136214	Avr1b-1 avirulence	2.0E-80
6.	jgi Physo1_1 136280	Avr1b-1 avirulence	1.8E-69
7.	jgi Physo1_1 136284	Avirulence 1b	4.4E-70
8.	jgi Physo1_1 139437	Avr1b-1 avirulence	1.4E-74
9.	jgi Physo1_1 139466	Avirulence 1b	1.9E-67
10.	jgi Physo1_1 139921	Avirulence 1b	1.1E-73
11.	jgi Physo1_1 139923	Avirulence 1b	1.1E-63
12.	jgi Physo1_1 142101	Avirulence 1b	4.9E-105
13.	jgi Physo1_1 142795	Avr1b-1 avirulence	1.7E-86
14.	jgi Physo1_1 131199	Avr1b-1 avirulence	1.4E-134
15.	jgi Physo1_1 135099	Avr1b-1 avirulence	1.8E-116
16.	jgi Physo1_1 140616	Avr1b-1 avirulence	2.1E-132

(2)

Table 2
Annotation results showing the occurrence of elicitors with Seq ID, functional description, and the corresponding E-value in (1) *P. ramorum* and (2) *P. sojae*.

S. N.	<i>Phytophthora ramorum</i> ID	Description	Length	E-value
1.	jgi Phyra1_1 71532	Elicitin-Like Mating M25	247	9.9E-179
2.	jgi Phyra1_1 71746	Elicitin RAL6	176	5.62E-90
3.	jgi Phyra1_1 76630	Elicitin RAL13C	137	2.37E-78
4.	jgi Phyra1_1 77586	Elicitin-Like Mating M25	251	0E0
5.	jgi Phyra1_1 78652	Elicitin Vex1	135	1.17E-79
6.	jgi Phyra1_1 96566	Elicitin RAL11D	138	4.9E-82

(1)

S.N.	<i>Phytophthora sojae</i> ID	Description	Length	E-value
1.	jgi Physo1_1 108257	Elicitin-Like Mating M25	242	2.74E-161
2.	jgi Physo1_1 134055	Elicitin SOL11C	136	2.56E-92
3.	jgi Physo1_1 135435	Elicitin Vex1	157	3.69E-98
4.	jgi Physo1_1 144033	Elicitin	135	1.07E-93
5.	jgi Phyra1_1 108901	Elicitin SOL6	172	4.77E-89

(2)

Moreover, SSPs (small secreted proteins) and small cysteine-rich protein have been reported and some of them were found to be common in the two species as depicted in the table 3. SSPs have been reported to be species-specific and could be potential targets to control the pathogenicity of the concerned species²⁹. Small-cysteine rich proteins are known fungal effectors that trigger resistance or

susceptibility in specific host plants³⁰. RXLR effector is known to be one of the most diverse groups of cytoplasmic effector families associated with oomycetes plant pathogen which has the potential to get delivered into the host cell and suppresses its immunity³¹. In this study, we reported species-specific RXLR effector proteins as mentioned in the table 4.

Table 3
Comparative view of small cysteine-rich proteins with Seq ID, functional description and the corresponding E-value in (a) *P. ramorum* and (b) *P. sojae*. First two proteins were predicted to be common in both the species.

S.N.	<i>Phytophthora ramorum</i> ID	Description	Length	E-value
1.	jgi Phyra1_1 43241	Small cysteine rich SCR91	73	7.6E-11
2.	jgi Phyra1_1 71697	Small cysteine rich SCR108	112	1.0E-42
3.	jgi Phyra1_1 78603	Small cysteine rich	145	2.6E-82

(a)

S.N.	<i>Phytophthora sojae</i> ID	Description	Length	E-value
1.	jgi Physo1_1 111409	Small cysteine rich SCR91	72	1.5E-31
2.	jgi Physo1_1 156264	Small cysteine rich SCR108	206	9.7E-131

(b)

Table 4
Comparative view: Proteins belonging to RXLR family with Seq ID, functional description and the corresponding E-value in (a) *P. ramorum* and (b) *P. sojae*.

S.N.	<i>Phytophthora ramorum</i> ID	Description	Length	E-value
1.	jgi Phyra1_1 78800	RXLR-class effector Avh247	127	4.3E-88
2.	jgi Phyra1_1 83417	RXLR-class effector Avh108	138	1.6E-6
3.	jgi Phyra1_1 79106	RXLR effector	122	7.2E-23
4.	jgi Phyra1_1 85154	RXLR effector	182	1.8E-5

(a)

S.N.	<i>Phytophthora sojae</i> ID	Description	Length	E-value
1.	jgi Physo1_1 139461	RXLR effector family	190	4.4E-139
2.	jgi Physo1_1 140615	RXLR effector Avh205	119	7.4E-82
3.	jgi Physo1_1 140618	RXLR effector Avh205	127	3.2E-88

(b)

In this study, we have also investigated the occurrence of effector proteins clusters in the two species as represented in figure 3. The result depicts the presence of 129 in-paralogs from *P. ramorum* and *P. sojae* each. Also, 102 groups of

orthologous have been predicted. The orthologous proteins clusters observed in the current study points towards the fact that *Phytophthora* spp. evolved through a remarkable degree of convergent evolution³².

Conclusion

Computational analysis and manual evaluation of the secretome of *Phytophthora* spp. could provide researchers with a comprehensive set of secretory proteins. Nevertheless, detailed studies and experimental validation of the predicted proteins will definitely enable an understanding of the molecular basis of host-pathogen interaction. Furthermore, identification of critical avirulence factors and elucidation of their functions promises to provide insight into plant defense mechanisms and to create a basis for the development of associated and improved strategies for the control of plant disease.

Most importantly, in this study we have reported host-specific effector proteins which are species-specific and lack similarity to known proteins which will prove to be potential target for pathogen control. With the current study, we would like to anticipate that the predicted data will provide a basis for the development of improved disease control strategies and promote research in this indispensable and complex area.

Acknowledgement

The authors would like to thank DBT (Sub-Distributed Information Centre) for funding.

References

- O'Hanlon R., Choiseul J., Corrigan M., Catarama T. and Destefanis M., Diversity and detection of *Phytophthora* species from trade and non-trade environments in Ireland, *EPPO Bull.*, **46**, 594–602 (2016)
- Delaunoy B., Jeandet P., Clément C., Baillieux F., Dorey S. and Cordelier S., Uncovering plant-pathogen crosstalk through apoplastic proteomic studies, *Front. Plant Sci.*, **5**, 249 (2014)
- Huitema E., Bos J.I., Tian M., Win J., Waugh M.E. and Kamoun S., Linking sequence to phenotype in *Phytophthora*-plant interactions, *Trends Microbiol.*, **12**(4), 193–200 (2004)
- Hadji Mollahosseini S., Host-pathogen interactions in root infecting oomycete species, Doctoral Dissertation, Swedish University of Agricultural Sciences, Uppsala (2015)
- Garbelotto M. and Hayden K.J., Sudden Oak Death: Interactions of the exotic oomycete *Phytophthora ramorum* with naive North American hosts, *Eukaryot. Cell*, **11**, 1313–1323 (2012)
- Rizzo D.M. and Garbelotto M., Sudden oak death: endangering California and Oregon forest ecosystems, *Frontiers Ecol. Environ.*, **1**(4), 197–204 (2003)
- Davidson J.M., Werres S., Garbelotto M., Hansen E.M. and Rizzo D.M., Sudden oak death and associated diseases caused by *Phytophthora ramorum*, *Plant Health Prog.*, doi:10.1094/PHP-2003-0707-01-DG (2003)
- Manter D.K., Kelsey R.G. and Karchesy J.J., Photosynthetic declines in *Phytophthora ramorum*-infected plants develop prior to water stress and in response to exogenous application of elicitors, *Phytopathology*, **97**(7), 850–856 (2007)
- Appiah A.A., Jennings P. and Turner J.A., *Phytophthora ramorum*: one pathogen and many diseases, an emerging threat to forest ecosystems and ornamental plant life, *Mycologist*, **18**(4), 145–150 (2004)
- Morris P.F., Bone E. and Tyler B.M., Chemotropic and contact responses of *Phytophthora sojae* hyphae to soybean isoflavonoids and artificial substrates, *Plant Physiol.*, **117**(4), 1171–1178 (1998)
- Tyler B.M., Tripathy S., Zhang X., Dehal P., Jiang R.H., Aerts A., Arredondo F.D., Baxter L., Bensasson D., Beynon J.L. and Chapman J., *Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis, *Science*, **313**, 1261–1266 (2006)
- Petersen T.N., Brunak S., von Heijne G. and Nielsen H., SignalP 4.0: Discriminating signal peptides from transmembrane regions, *Nat. Methods*, **8**, 785–786 (2011)
- Sonnhammer E.L., Von Heijne G. and Krogh A., A hidden Markov model for predicting transmembrane helices in protein sequences, *Ismb*, **6**, 175–182 (1998)
- Krogh A., Larsson B., Von Heijne G. and Sonnhammer E.L., Predicting transmembrane protein topology with a Hidden Markov Model: Application to complete genomes, *J. Mol. Biol.*, **305**, 567–580 (2001)
- Bendtsen J.D., Jensen L.J., Blom N., Von Heijne G. and Brunak S., Feature-based prediction of non-classical and leaderless protein secretion, *Protein Eng. Des. Sel.*, **17**, 349–356 (2004)
- Emanuelsson O., Nielsen H., Brunak S. and Von Heijne G., Predicting subcellular localization of proteins based on their N-terminal amino acid sequence, *J. Mol. Biol.*, **300**, 1005–1016 (2000)
- Fankhauser N. and Mäser P., Identification of GPI anchor attachment signals by a Kohonen self-organizing map, *Bioinformatics*, **21**, 1846–1852 (2005)
- Sperschneider J., Gardiner D.M., Dodds P.N., Tini F., Covarelli L., Singh K.B., Manners J.M. and Taylor J.M., Effector P: Predicting fungal effector proteins from secretomes using machine learning, *New Phytol.*, **210**, 743–761 (2016)
- Altschul S.F., Gish W., Miller W., Myers E.W. and Lipman D.J., Basic local alignment search tool, *J. Mol. Biol.*, **215**, 403–410 (1990)
- Conesa A., Götz S., García-Gómez J.M., Terol J., Talón M. and Robles M., Blast2GO: A universal tool for annotation, visualization and analysis in functional genomics research, *Bioinformatics*, **21**, 3674–3676 (2005)
- Östlund G., Schmitt T., Forslund K., Köstler T., Messina D.N., Roopra S., Frings O. and Sonnhammer E.L., In Paranoid 7: New algorithms and tools for eukaryotic orthology analysis, *Nucleic Acids Res.*, **38**, D196–D203 (2010)
- Guyon K., Balagué C., Roby D. and Raffaele S., Secretome analysis reveals effector candidates associated with broad host range necrotrophy in the fungal plant pathogen *Sclerotinia sclerotiorum*, *BMC Genomics*, **15**, 336 (2014)

23. Kwon M.S., Kim K., Lee S., Chung W., Yi S.G., Namkung J. and Park T., GWAS-GMDR: A program package for genome-wide scan of gene-gene interactions with covariate adjustment based on multifactor dimensionality reduction, *Bioinformatics and Biomedicine Workshops (BIBMW)*, IEEE International Conference, 703–707 (2011)
24. Rittenour W.R. and Harris S.D., Glycosylphosphatidylinositol-Anchored Proteins in *Fusarium graminearum*: Inventory, variability and virulence, *PLoS ONE*, **8**, e81603 (2013)
25. Sonah H., Deshmukh R.K. and Bélanger R.R., Computational prediction of effector proteins in fungi: Opportunities and challenges, *Front. Plant Sci.*, **7**, 126 (2016)
26. Narayanasamy P., Molecular Biology in Plant Pathogenesis and Disease Management, Disease Management, Springer Science & Business Media (2008)
27. Uhlíková H., Obořil M., Klemková J., Šedo O., Zdráhal Z., Kašparovský T., Skládal P. and Lochman J., Elicitin-induced distal systemic resistance in plants is mediated through the protein–protein interactions influenced by selected lysine residues, *Front. Plant Sci.*, **7**, 59 (2016)
28. Saubeau G., Gaillard F., Legentil L., Nugier-Chauvin C., Ferrières V., Andrivon D. and Val F., Identification of three elicitors and a galactan-based complex polysaccharide from a concentrated culture filtrate of *Phytophthora infestans* efficient against *Pectobacterium atrosepticum*, *Molecules*, **19**, 15374–15390 (2014)
29. Cheng Q., Wang H., Xu B., Zhu S., Hu L. and Huang M., Discovery of a novel small secreted protein family with conserved N-terminal IGY motif in Dikarya fungi, *BMC Genomics*, **15**(1), 1151 (2014)
30. Lu S. and Edwards M.C., Genome-wide analysis of small secreted cysteine-rich proteins identifies candidate effector proteins potentially involved in *Fusarium graminearum*–wheat interactions, *Phytopathology*, **106**(2), 166-176 (2016)
31. Ye W., Wang Y. and Wang Y., Bioinformatics analysis reveals abundant short alpha-helices as a common structural feature of oomycete RxLR effector proteins, *PloSOne*, **10**(8), e0135240 (2015)
32. Bailey B.A. and Meinhardt L.W., Cacao diseases: A history of old enemies and new encounters, Springer, 633 (2016).

(Received 18th March 2017, accepted 02nd May 2017)