

Erratum: Asymptotic and near-target direct breakup of ${}^6\text{Li}$ and ${}^7\text{Li}$ [Phys. Rev. C **93, 044605 (2016)]**

Sunil Kalkal, E. C. Simpson, D. H. Luong, K. J. Cook, M. Dasgupta, D. J. Hinde, I. P. Carter, D. Y. Jeung,
G. Mohanto, C. S. Palshetkar, E. Prasad, D. C. Rafferty, C. Simenel, K. Vo-Phuoc, E. Williams, L. R. Gasques,
P. R. S. Gomes, and R. Linares

(Received 24 May 2016; published 20 June 2016)

DOI: [10.1103/PhysRevC.93.069904](https://doi.org/10.1103/PhysRevC.93.069904)

The breakup cross sections in Appendix A had an error of 6–7% due to an error in the normalization procedure used to extract the cross sections. The correct cross sections are given in the table below. The conclusions of the article are not changed.

TABLE III. The asymptotic, near-target, and total direct breakup differential cross sections (in mb/sr) of ${}^{6,7}\text{Li}$ in interactions with the ${}^{209}\text{Bi}$ target at $E_{\text{lab}} = 29.0$ MeV.

$\theta_{\text{c.m.}}$ (deg)	${}^6\text{Li}$			${}^7\text{Li}$		
	Asymptotic	Near-target	Total	Asymptotic	Near-target	Total
122	0.79 ± 0.05	3.04 ± 0.09	3.84 ± 0.10	0.019 ± 0.003	0.29 ± 0.03	0.31 ± 0.03
126	0.88 ± 0.06	2.89 ± 0.08	3.77 ± 0.10	0.019 ± 0.002	0.30 ± 0.02	0.32 ± 0.02
130	0.92 ± 0.06	2.78 ± 0.06	3.69 ± 0.08	0.018 ± 0.002	0.27 ± 0.02	0.29 ± 0.02
134	0.94 ± 0.08	2.83 ± 0.05	3.77 ± 0.09	0.018 ± 0.002	0.28 ± 0.02	0.30 ± 0.02
138	0.94 ± 0.09	2.75 ± 0.07	3.69 ± 0.11	0.024 ± 0.003	0.30 ± 0.02	0.33 ± 0.02
142	1.00 ± 0.08	2.77 ± 0.08	3.77 ± 0.11	0.031 ± 0.003	0.30 ± 0.03	0.33 ± 0.03
146	0.99 ± 0.10	2.71 ± 0.07	3.70 ± 0.12	0.028 ± 0.003	0.32 ± 0.03	0.35 ± 0.03
150	1.04 ± 0.19	2.78 ± 0.08	3.82 ± 0.21	0.032 ± 0.004	0.34 ± 0.06	0.37 ± 0.06
154	1.10 ± 0.14	2.76 ± 0.10	3.86 ± 0.17	0.029 ± 0.005	0.33 ± 0.07	0.36 ± 0.07
158	1.12 ± 0.17	2.70 ± 0.09	3.82 ± 0.19	0.038 ± 0.007	0.33 ± 0.06	0.37 ± 0.06
162	1.01 ± 0.14	2.52 ± 0.08	3.53 ± 0.16	0.035 ± 0.006	0.32 ± 0.07	0.36 ± 0.07