

A Study on Isomorphic Properties of Circulant Graphs

V. Vilfred Kamalappan *

Department of Mathematics, Central University of Kerala Periyar, Kasaragod, Kerala, 671316 India

ARTICLE INFO

Article history:

Received: 13 October, 2017

Accepted: 06 November, 2017

Online: 20 December, 2017

Keywords:

Self-Complementary circulant graphs

Type-1 and Type-2 isomorphic circulant graphs

Cartesian product and factorization

prime and composite circulant graphs

Factorization theorem, Fundamental theorem

ABSTRACT

$C_n(R)$ denotes circulant graph $C_n(r_1, r_2, \dots, r_k)$ of order n for a set $R = \{r_1, r_2, \dots, r_k\}$ where $1 \leq r_1 < r_2 < \dots < r_k \leq \lfloor \frac{n}{2} \rfloor$. Circulant graph $C_n(R)$ is said to have the Cayley Isomorphism (CI) property if whenever $C_n(S)$ is isomorphic to $C_n(R)$, there is some $a \in \mathbb{Z}_n^*$ for which $S = aR$. In this paper, isomorphic properties of circulant graphs that includes (i) Self-complementary circulant graphs; (ii) Type-2 isomorphism, a new type of isomorphism other than already known Adam's isomorphism of circulant graphs and (iii) Cartesian product and factorization of circulant graphs similar to the theory of product and factorization of natural numbers are studied. New abelian groups are obtained from these isomorphic circulant graphs. Type-2 isomorphic circulant graphs have the property that they are isomorphic graphs without Cayley Isomorphism (CI) property and thereby new families.

1 Introduction

This paper is an extension of work originally presented in ICMSAO2017 [1] and covers the author's study on a few isomorphic properties of circulant graphs that includes (i) Existence of self-complementary circulant graphs; (ii) Type-2 isomorphism, a new type of isomorphism other than already known Adam's isomorphism of circulant graphs that helps to obtain graphs without CI-property and abelian groups and (iii) Cartesian product and factorization of circulant graphs similar to the theory of product and factorization of natural numbers.

Beauty comes out of symmetry as well as asymmetry. Investigation of symmetries/asymmetries of structures yield powerful results in Mathematics. Circulant graphs form a class of highly symmetric mathematical (graphical) structures. In 1846 Catalan (cf. [2]) introduced circulant matrices and properties of circulant graphs have been investigated by many authors [1-20]. An excellent account of circulant matrices can be found in the book by Davis [2] and circulant graphs in the article [11].

If a graph G is circulant, then its adjacency matrix $A(G)$ is circulant. It follows that if the first row of the adjacency matrix of a circulant graph is $[a_1, a_2, \dots, a_n]$, then $a_1 = 0$ and $a_i = a_{n-i+2}$, $2 \leq i \leq n$ [15, 17].



Through-out this paper, for a set $R = \{r_1, r_2, \dots, r_k\}$,

$C_n(R)$ denotes circulant graph $C_n(r_1, r_2, \dots, r_k)$ where $1 \leq r_1 < r_2 < \dots < r_k \leq \lfloor \frac{n}{2} \rfloor$. Only connected circulant graphs of finite order are considered, $V(C_n(R)) = \{v_0, v_1, v_2, \dots, v_{n-1}\}$ with v_i adjacent to v_{i+r} for each $r \in R$, subscript addition taken modulo n and all cycles have length at least 3, unless otherwise specified, $0 \leq i \leq n-1$. However when $\frac{n}{2} \in R$, edge $v_i v_{i+\frac{n}{2}}$ is taken as a single edge for considering the degree of the vertex v_i or $v_{i+\frac{n}{2}}$ and as a double edge while counting the number of edges or cycles in $C_n(R)$, $0 \leq i \leq n-1$. Generally, write C_n for $C_n(1)$ and $C_n(1, 2, \dots, \lfloor \frac{n}{2} \rfloor)$ for K_n . We will often assume, with-out further comment, that the vertices are the corners of a regular n -gon, labeled clockwise. Circulant graphs $C_{16}(1, 2, 7)$, $C_{16}(2, 3, 5)$, $C_{27}(1, 3, 8, 10)$, $C_{27}(3, 4, 5, 13)$ and $C_{27}(2, 3, 7, 11)$ are shown in Figures 1 - 5. Now, let us consider the following definitions and results that are useful in the subsequent sections.

Definition 1.1. [17] Let n and r be positive integers with $n \geq 4$ and $r < \frac{n}{2}$. Then, clearly, $C_n(r)$ consists of a collection of cycles $(v_0 v_r v_{2r} \dots v_0)$, $(v_1 v_{1+r} v_{1+2r} \dots v_1)$, \dots , $(v_{r-1} v_{2r-1} v_{3r-1} \dots v_{r-1})$. If $d = \gcd(n, r)$, then there are d such disjoint cycles, each of length $\frac{n}{d}$. We say that each of these cycles is of period r , length $\frac{n}{d}$ and rotation $\frac{r}{d}$.

If $r = \frac{n}{2}$, then obviously $C_n(r)$ is just a 1 - factor. Since $C_n(R)$ is just the union of the cycles $C_n(r)$ for $r \in R$, we have a decomposition of $C_n(R)$.

*Corresponding Author: V. Vilfred Kamalappan Email: vilfredkamalv@cukerala.ac.in, Contact No. 0091 9443577502



Theorem 1.2. [17] Let $r \in R$. Then, in $C_n(R)$, the length of a cycle of period r is $\frac{n}{\gcd(n,r)}$ and the number of disjoint periodic cycles of period r is $\gcd(n,r)$. \square

Corollary 1.3. [17] In $C_n(R)$, the length of a cycle of period r is n if and only if $\gcd(n,r) = 1, r \in R$. \square

Remark 1.4. [17] Let $|R| = k$. Then, the circulant graph $C_n(R)$ for a set $R = \{r_1, r_2, \dots, r_k\}$ is $(2k-1)$ -regular if $\frac{n}{2} \in R$ and $2k$ -regular otherwise. \square

The following Lemmas are useful to obtain one-to-one mappings.





Lemma 1.5. [15] Let A and B be two non-empty sets. Let $f : A \rightarrow B$ be a mapping. Then, f is one-to-one if and only if $f|_{A'}$ is one-to-one for every non-empty subset A' of A . \square

Lemma 1.6. [15] Let A and B be non-empty sets and A_1, A_2, \dots, A_k be a partition of A (each A_i being non-empty, $i = 1, 2, \dots, k$). Let $f : A \rightarrow B$ be a mapping. Then f is one-to-one if and only if $f|_{A_i}$ is one-to-one for every $i, i = 1, 2, \dots, k$. \square

2 On self-complementary circulant graphs

In 1962, Horst Sachs [14] proved that the sufficient condition for the existence of a self-complementary circulant graph of order n is that every prime factor p of n should satisfy $p \equiv 1 \pmod{4}$. He also conjectured that the sufficient condition is a necessary one. We have proved that the self-complementary circulant graph on n vertices does not exist if n has any prime factor which is not of the form $4m + 1, m \in \mathbb{N}$. Thereby, we establish that the sufficient condition is also a necessary one. The proof is based on counting the number of disjoint cycles of a particular length in K_n and is given in this section [4, 17]. Graphs given in Figures 6 and 7 are self-complementary but not of

circulant whereas graphs given in Figures 8 and 9 are self-complementary circulant graphs.



Theorem 2.1. [17] If $C_n(R) \cong C_n(S)$, then there is a bijection f from R to S so that for all $r \in R, \gcd(n,r) = \gcd(n, f(r))$.

Proof. The proof is by induction on the order of R . \square

Theorem 2.2. [9] If graph G of order n is self-complementary, then $n \equiv 0, 1 \pmod{4}$. \square

Theorem 2.3. [17] If $C_n(R)$ for a set $R = \{r_1, r_2, \dots, r_k\}$ is self-complementary, then $n \equiv 1 \pmod{4}$.

Proof. Using Theorem 2.2, we get $n \equiv 0, 1 \pmod{4}$. When n is even, a circulant graph is of even degree if and only if its complement is of odd degree since any circulant graph is a regular graph. Thus, self-complementary circulant graph of even order doesn't exist. Hence, we get the result. \square

Theorem 2.4. [17] If $C_n(R)$ for a set $R = \{r_1, r_2, \dots, r_k\}$ is self-complementary, then $n = 4m + 1, k = m$ and $|S_i|$ is even where $S_i = \{j : \gcd(n, j) = \gcd(n, i), 1 \leq j \leq \lfloor \frac{n}{2} \rfloor\}$, for all $i, i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$.

Proof. Using Theorem 2.3, we get, $n = 4m + 1$. This implies that the degree (of each vertex) of a self-complementary circulant graph of order $4m + 1$ is $2m$ which implies $k = m$.

Let $C_{4m+1}(R)$ be a self-complementary circulant graph for $R = \{r_1, r_2, \dots, r_m\}$. If it contains a cycle of period r (and of length $\frac{n}{\gcd(n,r)}$, using Theorem 1.2), then its complement also contains a periodic cycle of period, say, s such that $\frac{n}{\gcd(n,s)} = \frac{n}{\gcd(n,r)}$, $1 \leq r, s \leq \lfloor \frac{n}{2} \rfloor$, using Theorem 1.2. This implies that $\gcd(n,s) = \gcd(n,r)$. Here we consider that the cycles of periods r_i and $n - r_i$ are the same in $C_n(r_1, r_2, \dots, r_k), 1 \leq i \leq k$. Combining the above arguments, we get the result. \square

Remark 2.5. The above theorem states that if a self-complementary circulant graph of order n exists, then $n = 4m + 1$ and so it is $2m$ -regular and the number of periodic cycles of length i in K_n is always even for each i , $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$.

Theorem 2.6. [17] Self-complementary circulant graph of order n doesn't exist when n has any prime factor of the form $2(2m - 1) + 1$, $m \in \mathbb{N}$.

Proof. Using Theorem 2.3, $n = 4a + 1$, $a + 1 \in \mathbb{N}$.

Let $n = p_1^{n_1} p_2^{n_2} \dots p_j^{n_j}$ where p_1, p_2, \dots, p_j are the (odd) prime factors of n . Let $p_i \neq 4m + 1$ for at least one i , $1 \leq i \leq j$ and for any $m \in \mathbb{N}$. This implies, $p_i = 2(2q - 1) + 1$ for some $q \in \mathbb{N}$. Consider the circulant graph $C_{4a+1}(1, 2, \dots, 2a) \cong K_{4a+1} = K_n$. Let r be the natural number such that $r = \frac{n}{p_i}$.

Aim To find out all natural numbers lying between 1 and n such that g.c.d. of each one of them with n is exactly r .

Let $\gcd(n, pr + s) = r$ where $p, s + 1 \in \mathbb{N}$ such that $0 \leq s < r$ and $1 \leq pr + s \leq n = rp_i$. This implies that $s = 0$ and so $\gcd(n, rp + s) = \gcd(n, rp) = \gcd(rp_i, rp) = r$ where $1 \leq pr \leq p_i r = n$. Thus we get $\gcd(p, p_i) = 1$ where $p \leq p_i$ which is greater than 1. Therefore the possible values of p are $1, 2, \dots, p_i - 1$.

Thus $r, 2r, 3r, \dots, (p_i - 1)r$ are the only numbers lying between 1 and n such that g.c.d. of n with each one of them is exactly r . This implies $r, 2r, 3r, \dots, (p_i - 1)r$ are the possible periods of cycles of length p_i each, in $C_n(1, 2, \dots, \lfloor \frac{n}{2} \rfloor)$ since the length of a cycle of period rp in $C_n(1, 2, \dots, \lfloor \frac{n}{2} \rfloor)$ is $\frac{n}{\gcd(n, rp)} = \frac{rp_i}{\gcd(rp_i, rp)} = \frac{p_i}{\gcd(p_i, p)} = p_i$ for $p = 1, 2, \dots, p_i - 1$, using Theorem 1.2.

In $C_n(1, 2, \dots, \lfloor \frac{n}{2} \rfloor)$,

the cycles of period r and $n - r (= (p_i - 1)r)$ are the same,

the cycles of period $2r$ and $(p_i - 2)r$ are the same,

...

the cycles of period $\frac{(p_i - 1)r}{2}$ and $\frac{(p_i + 1)r}{2}$ are the same.

Thus, there are $\frac{p_i - 1}{2}$ number of possible distinct periodic cycles of (periods $r, 2r, \dots, \frac{p_i - 1}{2}r$ and) length p_i , each in $C_n(1, 2, \dots, \lfloor \frac{n}{2} \rfloor)$.

Now, $\frac{p_i - 1}{2} = 2q - 1$ is an odd number. This implies, any circulant graph of order n and its complementary circulant graph contain unequal number of periodic cycles of length p_i , each. This implies that self-complementary circulant graph of order n does not exist when n contains any prime factor of the form $2(2q - 1) + 1$, $q \in \mathbb{N}$, by Remark 2.5. \square

Thus, when $n = 9, 21, 33, 49, 57, 69, 77, 81, 93$, etc., self-complementary circulant graph doesn't exist on n vertices, by Theorem 2.6, even though in each case, $n \equiv 1 \pmod{4}$.

Now, by combining Theorem 2.6 and the sufficient condition for the existence of a self-complementary circulant graph of order n , we get the following result.

Theorem 2.7. [17] The necessary and sufficient condition for the existence of a self-complementary circulant

graph of order n is that each prime factor p of n should satisfy $p \equiv 1 \pmod{4}$. \square

3 On Isomorphism of Circulant Graphs

In this section, Type-2 isomorphism, a new type of isomorphism different from already known Adam's isomorphism of circulant graphs, main results related to it and families of new abelian groups obtained from isomorphic circulant graphs are presented. Type-2 isomorphic circulant graphs have the property that they are isomorphic graphs without Cayley Isomorphism (CI) property.

Definition 3.1. [12] A circulant graph $C_n(R)$ is said to have the CI-property if whenever $C_n(S)$ is isomorphic to $C_n(R)$, there is some $a \in \mathbb{Z}_n^*$ for which $S = aR$.

Lemma 3.2. [16] Let S be a non-empty subset of \mathbb{Z}_n and $x \in \mathbb{Z}_n$. Define a mapping $\Phi_{n,x} : S \rightarrow \mathbb{Z}_n$ such that $\Phi_{n,x}(s) = xs$ for every $s \in S$ under multiplication modulo n . Then, $\Phi_{n,x}$ is bijective if and only if $S = \mathbb{Z}_n$ and $\gcd(n, x) = 1$. \square

Definition 3.3. [3] Circulant graphs $C_n(R)$ and $C_n(S)$ for $R = \{r_1, r_2, \dots, r_k\}$ and $S = \{s_1, s_2, \dots, s_k\}$ are Adam's isomorphic if there exists a positive integer x relatively prime to n with $S = \{xr_1, xr_2, \dots, xr_k\}_n^*$ where $\langle r_i \rangle_n^*$ the reflexive modular reduction of a sequence $\langle r_i \rangle$ is the sequence obtained by reducing each r_i modulo n to yield r'_i and then replacing all resulting terms r'_i which are larger than $\frac{n}{2}$ by $n - r'_i$.

Lemma 3.4. [16] Let $m, r, t \in \mathbb{Z}_n \in \gcd(n, r) = m > 1$ and $0 \leq t \leq \frac{n}{m} - 1$. Then the mapping $\Theta_{n,r,t} : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ defined by $\Theta_{n,r,t}(x) = x + jtm$ for every $x \in \mathbb{Z}_n$ under arithmetic modulo n is bijective where $x = j + qm$, $0 \leq j \leq m - 1$, $0 \leq q \leq \frac{n}{m} - 1$ and $j, q \in \mathbb{Z}_n$. \square

Theorem 3.5. [16] Let $V(C_n(R)) = \{v_0, v_1, \dots, v_{n-1}\}$, $V(K_n) = \{u_0, u_1, \dots, u_{n-1}\}$, $r \in R$ and $\gcd(n, r) = m > 1$. Then the mapping $\Theta_{n,r,t} : V(C_n(R)) \rightarrow V(K_n)$ defined by $\Theta_{n,r,t}(v_x) = u_{x+jtm}$ and $\Theta_{n,r,t}((v_x, v_{x+s})) = (\Theta_{n,r,t}(v_x), \Theta_{n,r,t}(v_{x+s}))$ for every $x \in \mathbb{Z}_n$, $x = j + qm$, $0 \leq j \leq m - 1$, $0 \leq q, t \leq \frac{n}{m} - 1$ and $s \in R$, under subscript arithmetic modulo n , for a set $R = \{r_1, r_2, \dots, r_k, n - r_k, n - r_{k-1}, \dots, n - r_1\}$ is one-to-one, preserves adjacency and $\Theta_{n,r,t}(C_n(R)) \cong C_n(R)$ for $t = 0, 1, 2, \dots, \frac{n}{m} - 1$. \square

Definition 3.6. [16] For a given $C_n(R)$ and for a particular value of t , $0 \leq t \leq \frac{n}{m} - 1$, if $\Theta_{n,r,t}(C_n(R)) = C_n(S)$ for some $S \subseteq [1, \frac{n}{2}]$ and $S \neq xR$ for all $x \in \Phi_n$ under reflexive modulo n , then $C_n(R)$ and $C_n(S)$ are called Type-2 isomorphic circulant graphs w.r.t. $r, r \in R$. In this case, subsets R and S of \mathbb{Z}_n are called Type-2 isomorphic subsets of \mathbb{Z}_n w.r.t. r .

Clearly, Type-2 isomorphic circulant graphs are circulant graphs without CI-property. We obtained the following results on Type-2 isomorphism.

Theorem 3.7. [16] For $n \geq 2, k \geq 3, 1 \leq 2s-1 \leq 2n-1, n \neq 2s-1, R = \{2s-1, 4n-2s+1, 2p_1, 2p_2, \dots, 2p_{k-2}\}$ and $S = \{2n - (2s - 1), 2n + 2s - 1, 2p_1, 2p_2, \dots, 2p_{k-2}\}$, circulant graphs $C_{8n}(R)$ and $C_{8n}(S)$ are Type-2 isomorphic (and without CI - property) where $\gcd(p_1, p_2, \dots, p_{k-2}) = 1$ and $n, s, p_1, p_2, \dots, p_{k-2} \in \mathbb{N}$. \square

Theorem 3.8. [16] For $R = \{2r-1, 2s-1, 2p_1, 2p_2, \dots, 2p_{k-2}\}, n \geq 2, k \geq 3, 1 \leq t \leq \lfloor \frac{n}{2} \rfloor, 1 \leq 2r-1 < 2s-1 \leq \lfloor \frac{n}{2} \rfloor, \gcd(p_1, p_2, \dots, p_{k-2}) = 1$ and $n, r, s, t, p_1, p_2, \dots, p_{k-2} \in \mathbb{N}$, if $\Theta_{n,2,t}(C_n(R))$ and $C_n(R)$ are Type-2 isomorphic circulant graphs for some t , then $n \equiv 0 \pmod{8}, 2r-1+2s-1 = \frac{n}{2}, t = \frac{n}{8}$ or $\frac{3n}{8}, 2r-1 \neq \frac{n}{8}, 1 \leq 2r-1 \leq \frac{n}{4}$ and $n \geq 16$. \square

Definition 3.9. [1] Let $Ad_n = \{\Phi_{n,x} : x \in \Phi_n\}, Ad_n(R) = \{\Phi_{n,x}(R) : x \in \Phi_n\} = \{xR : x \in \Phi_n\}$ and $Ad_n(C_n(R)) = T1_n(C_n(R)) = \{\Phi_{n,x}(C_n(R)) : x \in \Phi_n\} = \{C_n(xR) : x \in \Phi_n\}$ for a set $R = \{r_1, r_2, \dots, r_k, n-r_k, n-r_{k-1}, \dots, n-r_1\} \subseteq \mathbb{Z}_n$. Define 'o' in $Ad_n(C_n(R))$ such that $\Phi_{n,x}(C_n(R)) \circ \Phi_{n,y}(C_n(R)) = \Phi_{n,xy}(C_n(R))$ and $C_n(xR) \circ C_n(yR) = C_n((xy)R)$ for every $x, y \in \Phi_n$, under arithmetic modulo n . Clearly, $Ad_n(C_n(R))$ is the set of all circulant graphs that are Adam's isomorphic to $C_n(R)$ and $(Ad_n(C_n(R)), \circ) = (T1_n(C_n(R)), \circ)$ is an abelian group and we call it as the Adam's group or Type-1 group on $C_n(R)$ under 'o'.

Definition 3.10. [1] Let $V(C_n(R)) = \{v_0, v_1, \dots, v_{n-1}\}, V(K_n) = \{u_0, u_1, \dots, u_{n-1}\}, r \in R, m, q, t, t', x \in \mathbb{Z}_n$ such that $\gcd(n, r) = m > 1, x = j + qm, 0 \leq j \leq m-1$ and $0 \leq q, t, t' \leq \frac{n}{m} - 1$. Define $\Theta_{n,r,t} : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ and $\Theta_{n,r,t} : V(C_n(R)) \rightarrow V(K_n)$ such that $\Theta_{n,r,t}(x) = x + jtm, \Theta_{n,r,t}(v_x) = u_{x+jtm}$ and $\Theta_{n,r,t}((v_x, v_{x+s})) = (\Theta_{n,r,t}(v_x), \Theta_{n,r,t}(v_{x+s}))$ for every $x \in \mathbb{Z}_n$ and $s \in R$, under subscript arithmetic modulo n . Let $s \in \mathbb{Z}_n, V_{n,r} = \{\Theta_{n,r,t} : t = 0, 1, \dots, \frac{n}{m} - 1\}, V_{n,r}(s) = \{\Theta_{n,r,t}(s) : t = 0, 1, \dots, \frac{n}{m} - 1\}$ and $V_{n,r}(C_n(R)) = \{\Theta_{n,r,t}(C_n(R)) : t = 0, 1, \dots, \frac{n}{m} - 1\}$. Define 'o' in $V_{n,r}$ such that $\Theta_{n,r,t} \circ \Theta_{n,r,t'} = \Theta_{n,r,t+t'}, (\Theta_{n,r,t} \circ \Theta_{n,r,t'})(x) = \Theta_{n,r,t}(\Theta_{n,r,t'}(x)) = \Theta_{n,r,t}(x + jt'm) = (x + jt'm) + jtm = x + j(t+t')m = \Theta_{n,r,t+t'}(x)$ and $\Theta_{n,r,t}(C_n(R)) \circ \Theta_{n,r,t'}(C_n(R)) = \Theta_{n,r,t+t'}(C_n(R))$ for every $\Theta_{n,r,t}, \Theta_{n,r,t'} \in V_{n,r}$ where $t+t'$ is calculated under addition modulo $\frac{n}{m}$. Clearly, $(V_{n,r}(s), \circ)$ and $(V_{n,r}(C_n(R)), \circ)$ are abelian groups for every $s \in \mathbb{Z}_n$.

$V_{n,r}(C_n(R))$ contains all isomorphic circulant graphs of Type-2 of $C_n(R)$ w.r.t. r , if exist. Let $T2_{n,r}(C_n(R)) = \{C_n(R)\} \cup \{C_n(S) : C_n(S) \text{ is Type-2 isomorphic to } C_n(R) \text{ w.r.t. } r\}$. Thus, $T2_{n,r}(C_n(R)) = \{C_n(R)\} \cup \{\Theta_{n,r,t}(C_n(R)) : \Theta_{n,r,t}(C_n(R)) = C_n(S) \text{ and } C_n(S) \text{ is Type-2 isomorphic to } C_n(R) \text{ w.r.t. } r, 0 \leq t \leq \frac{n}{m} - 1\} = \{\Theta_{n,r,0}(C_n(R))\} \cup \{\Theta_{n,r,t}(C_n(R)) : \Theta_{n,r,t}(C_n(R)) = C_n(S) \text{ and } C_n(S) \text{ is Type-2 isomorphic to } C_n(R), 0 \leq t \leq \frac{n}{m} - 1\} \subseteq V_{n,r}(C_n(R))$ and $(T2_{n,r}(C_n(R)), \circ)$ is a subgroup of $(V_{n,r}(C_n(R)), \circ)$. Clearly, $T1_n(C_n(R)) \cap T2_{n,r}(C_n(R)) = \{C_n(R)\}$. And $C_n(R)$ has Type-2 isomorphic circulant graph w.r.t. r if and only if $T2_{n,r}(C_n(R)) \neq \{C_n(R)\}$ if and only if $T2_{n,r}(C_n(R)) \cap \{C_n(R)\} \neq \Phi$ if and only if $|T2_{n,r}(C_n(R))| > 1$ [1].

Definition 3.11. [1] For any circulant graph $C_n(R)$, if $T2_{n,r}(C_n(R)) \neq \{C_n(R)\}$, then $(T2_{n,r}(C_n(R)), \circ)$ is called the Type-2 group of $C_n(R)$ w.r.t. r under 'o'.

Theorem 3.12. [1] Let p be an odd prime and $k \geq 3$. Then, for $i = 1$ to $p, d_i = (i-1)np + 1$ and $R_i = \{d_i, np^2 - d_i, np^2 + d_i, 2np^2 - d_i, 2np^2 + d_i, 3np^2 - d_i, 3np^2 + d_i, \dots, (p-1)np^2 - d_i, (p-1)np^2 + d_i, np^3 - d_i, pp_1, pp_2, \dots, pp_{k-2}, p(np^3 - p_{k-2}), p(np^3 - p_{k-3}), \dots, p(np^3 - p_1)\}$, circulant graphs $C_{np^3}(R_i)$ are Type-2 isomorphic (and without CI-property) where $\gcd(p_1, p_2, \dots, p_{k-2}) = 1$ and $n, p_1, p_2, \dots, p_{k-2} \in \mathbb{N}$. \square

Theorem 3.13. [1] Let p be an odd prime, $k \geq 3, 1 \leq i \leq p, d_i = (i-1)np + 1, R_i = \{d_i, np^2 - d_i, np^2 + d_i, 2np^2 - d_i, 2np^2 + d_i, 3np^2 - d_i, 3np^2 + d_i, \dots, (p-1)np^2 - d_i, (p-1)np^2 + d_i, np^3 - d_i, pp_1, pp_2, \dots, pp_{k-2}, p(np^3 - p_{k-2}), p(np^3 - p_{k-3}), \dots, p(np^3 - p_1)\}$, $T2(R_i) = \{\Theta_{np^3,p,jn}(R_i) : j = 1, 2, \dots, p\}, T2(C_{np^3}(R_i)) = \{\Theta_{np^3,p,jn}(C_{np^3}(R_i)) : j = 1, 2, \dots, p\}, \gcd(p_1, p_2, \dots, p_{k-2}) = 1$ and $n, p_1, p_2, \dots, p_{k-2} \in \mathbb{N}$. Then $T2_{np^3,p}(R_i) = T2(R_j), T2_{np^3,p}(C_{np^3}(R_i)) = T2(C_{np^3}(R_j))$ and $(V_{np^3,p}(R_i), \circ), (V_{np^3,p}(C_{np^3}(R_i)), \circ)$ and $(T2_{np^3,p}(R_i), \circ)$ are abelian groups, $1 \leq i, j \leq p$. Moreover, $(T2_{np^3,p}(C_{np^3}(R_i)), \circ)$ is the Type-2 group of order p on $C_{np^3}(R_i)$ w.r.t. $r = p, 1 \leq i, j \leq p$. \square

Circulant graphs $C_{16}(1,2,7)$ and $C_{16}(2,3,5)$ are Type-2 isomorphic and $C_{27}(1,3,8,10), C_{27}(3,4,5,13)$ and $C_{27}(2,3,7,11)$ are also. See Figures 1-5.

4 Cartesian Product and Factorization of Circulant Graphs

Just as integers can be factored into prime numbers, there are many results on decompositions of structures throughout mathematics [6]. The standard products - Cartesian, lexicographic, tensor, and strong - all belong to a class of products introduced by Imrich and Izbicki and called B -products [8]. In this section, a few important results that are obtained in our study of Cartesian product and factorization of circulant graphs, similar to the theory of product and factorization of natural numbers, are presented (For details see [15]). Graphs $C_5 \square C_6$ and $C_{30}(5,6)$ are isomorphic and are given in Figures 12 and 13. One can see the difficulty of this study from this example.

Definition 4.1. [9] The cross product or Cartesian product of two simple graphs $G(V, E)$ and $H(W, F)$ is the simple graph $G \square H$ with vertex set $V \times W$ in which two vertices $u = (u_1, u_2)$ and $v = (v_1, v_2)$ are adjacent if and only if either $u_1 = v_1$ and $u_2 v_2 \in F$ or $u_2 = v_2$ and $u_1 v_1 \in E$.

Theorem 4.2. [15] Let G be a connected graph of order $n, n > 2$. Then, $P_2 \square G$ is circulant if and only if $G \cong H$ or $P_2 \square H$ where H is a connected circulant graph of odd order. \square

Theorem 4.3. [15] Let G be a connected graph of order $n \geq 2$. Then $C_4 \square G$ is circulant if and only if G is circulant of odd order. \square

Theorem 4.4. [15] If G and H are connected graphs and $G \square H$ is circulant, then G and H are circulants. \square

Graphs $P_2 \square C_3$ and $P_2 \square C_4$ are given in Figures 10 and 11 and $C_5 \square C_6$ and $C_{30}(5,6) \cong C_5 \square C_6$ in 12 and 13, respectively.




Fig.10. $P_2 \times C_3$




Fig.11. $C_6(2,3)$




Fig.12. $C_5 \times C_6$




Fig.13. $C_{30}(5,6) \cong C_5 \times C_6$

Theorem 4.5. [15] Let G and H be connected graphs, each of order > 2 . Then $G \square H$ is circulant if and only if G and H are circulants and satisfy one of the following conditions:

- (i) $G \cong C_m(R)$; $H \cong C_n(S)$ and $\gcd(m, n) = 1$.

- (ii) $G \cong C_{2m+1}(R)$; $H \cong C_{2n+1}(S)$, $P_2 \square C_{2n+1}(S)$, $C_4 \square C_{2n+1}(S)$ or $C_{2^k(2n+1)}(S)$ and $\gcd(2m+1, 2^k(2n+1)) = 1$, $k \in \mathbb{N}$.
- (iii) $G \cong P_2 \square C_{2m+1}(R)$; $H \cong C_{2n+1}(S)$ or $P_2 \square C_{2n+1}(S)$ and $\gcd(2m+1, 2n+1) = 1$.
- (iv) $G \cong C_{2^k(2m+1)}(R) \neq P_2 \square C_{2^{k-1}(2m+1)}(T)$ for any $C_{2^{k-1}(2m+1)}(T)$; $H \cong C_{2n+1}(S)$ and $\gcd(2^k(2m+1), 2n+1) = 1$, $k \in \mathbb{N}$.
- (v) $G \cong C_4 \square C_{2m+1}(R)$; $H \cong C_{2n+1}(S)$ and $\gcd(2m+1, 2n+1) = 1$.
- (vi) $G \cong C_{2^k(2m+1)}(R) \neq C_4 \square C_{2^{k-2}(2m+1)}(T)$, $P_2 \square C_{2^{k-1}(2m+1)}(U)$ for any $C_{2^{k-2}(2m+1)}(T)$ and $C_{2^{k-1}(2m+1)}(U)$; $H \cong C_{2n+1}(S)$ and $\gcd(2^k(2m+1), 2n+1) = 1$, $k \in \mathbb{N}$, $k \geq 2$. \square

Definition 4.6. [15] A non-trivial graph G is said to be prime if $G = G_1 \square G_2$ implies G_1 or G_2 is trivial; G is composite if it is not prime.

Definition 4.7. [15] If $C_m(R)$, $C_n(S)$ and $C_{mn}(T)$ are circulant graphs such that $C_m(R) \square C_n(S) \cong C_{mn}(T)$, then we say that $C_m(R)$ and $C_n(S)$ are divisors or factors of $C_{mn}(T)$.

Thus for any connected circulant graph, the graph and $C_1(\cdot) = K_1$ are always divisors and so we call them as *improper divisors* of the circulant graph. Divisors which are integer multiple of improper divisors also be called as improper divisors of the circulant graph. This doesn't arise since we consider divisors of connected graphs only. Divisor(s) other than improper divisors is called *proper divisor(s)* of the circulant graph.

Definition 4.8. [15] A circulant graph whose only divisors are improper is called a *prime circulant graph*. Other circulant graphs are called *composite circulant graphs*.

Theorem 4.9. [15] [Factorization Theorem On Circulant Graphs]

Let m and n be relatively prime integers. If $R \subseteq [1, \frac{m}{2}]$, $S \subseteq [1, \frac{n}{2}]$ and $T \subseteq [1, \frac{mn}{2}]$ with $T = d n R \cup d m S$ for some d such that $\gcd(mn, d) = 1$, then $C_{mn}(T) \cong C_m(R) \square C_n(S)$. \square

Theorem 4.10. [15] If $n \neq 4$ and $1 \in R$, then $C_n(R)$ is a prime circulant. \square

Corollary 4.11. [15] If $n \neq 4$ and R contains an integer relatively prime to n , then $C_n(R)$ is prime circulant. \square

Corollary 4.12. [15] If n is a prime power other than 4 and $C_n(R)$ is connected, then $C_n(R)$ is prime circulant for all $R \neq \emptyset$. \square

Theorem 4.13. [15] [Fundamental Theorem of Circulant Graphs]

Every connected circulant graph is the unique product of prime circulant graphs (uniqueness up to isomorphism). \square

Remark 4.14. [13, 15] If G is a connected graph such that $G \cong G_1 \square G_2 \square \dots \square G_k$, then the diameter of G , $\text{dia}(G) = \sum_{i=1}^k \text{dia}(G_i)$.

Thus, we can find the diameter of any given circulant graph, provided diameters of its prime circulant graphs are known. Also the above relation helps to generate (circulant) graphs of bigger diameters.

Remark 4.15. [15]

1. *In prime factorization of connected circulants $C_1() = K_1$ and $C_2 = P_2$ act similar to 1 and 2 among the set of all natural numbers, respectively. Thus, $C_1()$ is a unit, like 1 in number theory.*
2. *There exist two types of prime circulant graphs of order n , one with periodic cycle(s) of length n and the other without periodic cycle of length n .*
3. *The theory of factorization of circulants is similar to the theory of factorization of natural numbers and one of the very few well-known mathematical structures so vividly classified (expressed) in terms of prime factors. It can be applied in cryptography.*
4. *We developed VB programs POLY215.exe and POLY315.exe to show visually how the transformations $\Theta_{n,r,t}$ and $\Phi_{m,n}$ act on C_n and $C_m \square C_n$, respectively for different values of m and n , $m, n \in \mathbb{N}$.*
5. *An interesting problem is, for a given integer n , finding the number of prime (composite) circulant graphs of order either equal to n or less than or equal to n .*
6. *One can develop theories similar to the theory of Cartesian product and factorization of circulant graphs to the other standard products of circulant graphs.*

Conclusion This study covers a few isomorphic properties of circulant graphs. One can go for a similar study on Cayley graphs.

Acknowledgment The author expresses his sincere thanks to Prof. Lowell W Beineke, Indiana-Purdue University, U.S.A., Prof. Brian Alspach, University of Newcastle, Australia, Prof. M.I. Jinnah, University of Kerala, Trivandrum, India and Prof. V. Mohan, Thiyagarayar College of Engineering, Madurai, Tamil Nadu, India for their valuable suggestions and guidance and Dr. A. Christopher, Dr. P. Wilson and Mr. R. Sathesh of S.T. Hindu College, Nagercoil, India for their assistance to develop the VB programs to show how the transformations defined on circulant graphs to obtain their isomorphic graphs is taking place. He also expresses his gratitude to Central University of Kerala (CU Kerala), Periyar - 671 316, Kasaragod, Kerala, India and St. Jude's College, Thoothoor - 629 176,

Kanyakumari District, Tamil Nadu, India for providing facilities and CU Kerala and Lerroy Wilson Foundation, India (www.WillFoundation.co.in) for providing financial assistance to do this research work.

References

1. V. Vilfred, "A few properties of circulant graphs: Self-complementary, isomorphism, Cartesian product and factorization" in 7th International Conference on Modeling, Simulation and Applied Optimization (ICMSAO2017), American University of Sharjah, Sharjah, April 2017.
2. P. J. Davis, *Circulant Matrices*, Wiley, New York, 1979.
3. A. Adam, "Research problem 2-10", *J. Combinatorial Theory*, 3, 393, 1967.
4. B. Alspach, J. Morris and V. Vilfred, "Self-complementary circulant graphs", *Ars Com.*, 53, 187-191, 1999.
5. B. Alspach and T. B. Parsons, "Isomorphism of circulant graphs and digraphs", *Discrete Math.*, 25, 97-108, 1979.
6. R. Arratia, A. D. Barbour and Simon Tavaré, "Random Combinatorial Structures and Prime Factorizations", *A.M.S. Notices*, 44, 903-910, 1997.
7. F. T. Boesch and R. Tindell, "Circulant and their connectivities", *J. Graph Theory*, 8, 487-499, 1984.
8. B. Elspas and J. Turner, "Graphs with circulant adjacency matrices", *J. Combinatorial Theory*, 9, 297-307, 1970.
9. F. Harary, *Graph Theory*, Addison-Wesley, Reading, M.A., 1969.
10. W. Imrich and H. Izbicki, "Associative products of graphs", *Monatsh. Math.*, 80, 277-281, 1975.
11. I. Kra and S. Simanca, "On Circulant Matrices", *AMS Notices*, 59 (3), 368-377, 2012.
12. C. H. Li, "On isomorphisms of finite Cayley graphs-a survey", *Discrete Math.* 256, 301-334, 2002.
13. G. Sabidussi, "Graph multiplication", *Math. Z.*, 72, 446-457, 1960.
14. H. Sachs, "Über selbstkomplementäre Graphen", *Publ. Math. Debrecen*, 9, 270-288, 1962.
15. V. Vilfred, "A Theory of Cartesian Product and Factorization of Circulant Graphs", *Hindawi Pub. Corp. - J. Discrete Math.*, Vol. 2013, Article ID 163740, 10 pages.
16. V. Vilfred, "New Abelian Groups from Isomorphism of Circulant Graphs", *Proce. of Inter. Conf. on Applied Math. and Theoretical Computer Science*, St. Xaviers Catholic Engineering College, Nagercoil, Tamil Nadu, India, xiii-xvi, 2013.
17. V. Vilfred, " Σ -labelled Graphs and Circulant Graphs", *Ph.D. Thesis*, University of Kerala, Trivandrum, India, 1994.
18. V. Vilfred and P. Wilson, "New Families of Circulant graphs without Cayley Isomorphism Property with $m_i = 5$ ", *Int. Journal of Scientific and Innovative Mathematical Research*, 3 (6), 39-47, 2015.
19. V. Vilfred and P. Wilson, "New Family of Circulant Graphs without Cayley Isomorphism Property with $m_i = 7$ ", *IOSR Journal of Mathematics*, 12, 32-37, 2016.
20. H. Zhang, "Self-complementary symmetric graphs", *J. Graph Theory*, 16, 1-5, 1992.