



NAGESWARA RAO K.¹, GERMINA K.A.², SHAINI P.¹

ON THE DIMENSION OF VERTEX LABELING OF k -UNIFORM DCSL OF k -UNIFORM CATERPILLAR

A distance compatible set labeling (dcsL) of a connected graph G is an injective set assignment $f : V(G) \rightarrow 2^X$, X being a nonempty ground set, such that the corresponding induced function $f^\oplus : E(G) \rightarrow 2^X \setminus \{\emptyset\}$ given by $f^\oplus(uv) = f(u) \oplus f(v)$ satisfies $|f^\oplus(uv)| = k_{(u,v)}^f d_G(u, v)$ for every pair of distinct vertices $u, v \in V(G)$, where $d_G(u, v)$ denotes the path distance between u and v and $k_{(u,v)}^f$ is a constant, not necessarily an integer. A dcsL f of G is k -uniform if all the constant of proportionality with respect to f are equal to k , and if G admits such a dcsL then G is called a k -uniform dcsL graph. The k -uniform dcsL index of a graph G , denoted by $\delta_k(G)$ is the minimum of the cardinalities of X , as X varies over all k -uniform dcsL-sets of G . A linear extension L of a partial order $P = (P, \preceq)$ is a linear order on the elements of P , such that $x \preceq y$ in P implies $x \leq y$ in L , for all $x, y \in P$. The dimension of a poset P , denoted by $\dim(P)$, is the minimum number of linear extensions on P whose intersection is ' \preceq '. In this paper we prove that $\dim(\mathcal{F}) \leq \delta_k(P_n^{+k})$, where \mathcal{F} is the range of a k -uniform dcsL of the k -uniform caterpillar, denoted by P_n^{+k} ($n \geq 1, k \geq 1$) on ' $n(k+1)$ ' vertices.

Key words and phrases: k -uniform dcsL index, dimension of a poset, lattice.

¹ Department of Mathematics, Central University of Kerala, Kasaragod, Kerala 671314, India

² Department of Mathematics, University of Botswana, 4775 Notwane Rd., Private Bag UB 0022, Gaborone, Botswana
E-mail: karreynageswararao@gmail.com (Nageswara Rao K.), srgerminaka@gmail.com (Germina K.A.),
shainipv@gmail.com (Shaini P.)

INTRODUCTION

Acharya [1] introduced the notion of vertex *set-valuation* as a set-analogue of number valuation. For a graph $G = (V, E)$ and a nonempty set X , Acharya defined a *set-valuation* of G as an injective *set-valued* function $f : V(G) \rightarrow 2^X$, and defined a *set-indexer* $f^\oplus : E(G) \rightarrow 2^X \setminus \{\emptyset\}$ as a *set-valuation* such that the function given by $f^\oplus(uv) = f(u) \oplus f(v)$ for every $uv \in E(G)$ is also injective, where 2^X is the set of all subsets of X and ' \oplus ' is the binary operation of taking the symmetric difference of subsets of X .

Acharya and Germina [2], introduced the particular kind of set-valuation for which a metric, especially the cardinality of the symmetric difference, associated with each pair of vertices is k (where k be a constant) times that of the distance between them in the graph [2]. In other words, determine those graphs $G = (V, E)$ that admit an injective set-valued function $f : V(G) \rightarrow 2^X$, where 2^X is the power set of a nonempty set X , such that, for each pair of distinct vertices u and v in G , the cardinality of the symmetric difference $f(u) \oplus f(v)$ is k times

YΔK 519.1

2010 *Mathematics Subject Classification:* 05C78, 05C60.

The authors are thankful to the Department of Science and Technology, Government of India, New Delhi, for the financial support concerning the Major Research Project (Ref: No. SR/S4/MS : 760/12).