

Six unified results for reducibility of the Srivastava's triple hypergeometric series HA.

Source: Journal of Nonlinear Sciences & Applications (JNSA) . 2017, Vol. 10 Issue 6, p3038-3045. 8p.

Author(s): Junesang Choi; Rathie, Arjun K.

Abstract:

We aim to provide six unified results for reducibility of the Srivastava's triple hypergeometric series H_{A} . The results are obtained with the help of generalizations of classical summation theorems due to Kummer, Gauss second and Bailey for the series $F_{2}F_{1}$ which have recently been published. Our main findings are also shown to be specialized to yield several known results.

Copyright of Journal of Nonlinear Sciences & Applications (JNSA) is the property of Journal of Nonlinear Sciences & its Application (JNSA) and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use. This abstract may be abridged. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material for the full abstract.

For access to this entire article and additional high quality information, please check with your college/university library, local public library, or affiliated institution.



Important User Information: Remote access to EBSCO's databases is permitted to patrons of subscribing institutions accessing from remote locations for personal, non-commercial use. However, remote access to EBSCO's databases from non-subscribing institutions is not allowed if the purpose of the use is for commercial gain through cost reduction or avoidance for a non-subscribing institution.

[Privacy Policy](#) | [A/B Testing](#) | [Terms of Use](#) | [Copyright](#) | [Cookie Policy](#)

© 2021 EBSCO Industries, Inc. All rights reserved.