



NOTE ON THE EXTENDED WATSON'S SUMMATION THEOREM FOR THE SERIES ${}_4F_3(1)$

Junesang Choi*, Vivek Rohira and Arjun K. Rathie

Department of Mathematics
Dong University
Gyeongju 38066, Republic of Korea
e-mail: junesang@mail.dongguk.ac.kr

Department of Mathematics
Career Point University
Kota-325003, Rajasthan State
India
e-mail: vivekrohira@yahoo.com

Department of Mathematics
School of Physical Sciences
Central University of Kerala
Periye P.O., Kasaragod-671316
Kerala, India
e-mail: akrathie@cukerala.ac.in

Abstract

We aim to provide a new proof of the extended Watson's summation theorem for the series ${}_4F_3(1)$ due recently to Kim et al. [4]. Several interesting special cases of the extended Watson's summation theorem

Received: August 25, 2017; Revised: September 19, 2017; Accepted: October 13, 2017

2010 Mathematics Subject Classification: 33C20.

Keywords and phrases: generalized hypergeometric function, Watson summation theorem.

*Corresponding author

are also considered. We note that the method of proof used here is potentially useful in getting some other summation formulas for the series ${}_pF_q$.

1. Introduction

Throughout this paper, let ${}_pF_q$ denote the generalized hypergeometric series (see, for details, e.g., [1, 6], [7, Section 1.5]). Kim et al. [4] established extensions of various classical summation theorems for the series ${}_2F_1$, ${}_3F_2$ and ${}_4F_3$, two of which, for our present investigation, are recalled:

$$\begin{aligned}
 {}_3F_2 \left[\begin{matrix} a, b, d+1; \\ \frac{1}{2}(a+b+3), d; \end{matrix} \frac{1}{2} \right] &= \frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}a + \frac{1}{2}b + \frac{3}{2}\right)\Gamma\left(\frac{1}{2}a - \frac{1}{2}b - \frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}a - \frac{1}{2}b - \frac{3}{2}\right)} \\
 &\quad \times \left\{ \frac{\frac{1}{2}(a+b-1) - \frac{ab}{d}}{\Gamma\left(\frac{1}{2}a + \frac{1}{2}\right)\Gamma\left(\frac{1}{2}b + \frac{1}{2}\right)} + \frac{\frac{a+b+1}{d} - 2}{\Gamma\left(\frac{1}{2}a\right)\Gamma\left(\frac{1}{2}b\right)} \right\}
 \end{aligned} \tag{1.1}$$

and

$$\begin{aligned}
 &{}_4F_3 \left[\begin{matrix} a, b, c, d+1; \\ \frac{1}{2}(a+b+3), 2c, d; \end{matrix} 1 \right] \\
 &= \frac{2^{a+b-2}\Gamma\left(c + \frac{1}{2}\right)\Gamma\left(\frac{1}{2}a + \frac{1}{2}b + \frac{3}{2}\right)\Gamma\left(c - \frac{1}{2}a - \frac{1}{2}b - \frac{1}{2}\right)}{(a-b-1)(a-b+1)\Gamma\left(\frac{1}{2}\right)\Gamma(a)\Gamma(b)} \\
 &\quad \times \left\{ c_1 \frac{\Gamma\left(\frac{1}{2}a\right)\Gamma\left(\frac{1}{2}b\right)}{\Gamma\left(c - \frac{1}{2}a + \frac{1}{2}\right)\Gamma\left(c - \frac{1}{2}b + \frac{1}{2}\right)} + c_2 \frac{\Gamma\left(\frac{1}{2}a + \frac{1}{2}\right)\Gamma\left(\frac{1}{2}b + \frac{1}{2}\right)}{\Gamma\left(c - \frac{1}{2}a\right)\Gamma\left(c - \frac{1}{2}b\right)} \right\},
 \end{aligned} \tag{1.2}$$

provided $d \in \mathbb{C} \setminus \mathbb{Z}_0^-$, $\Re(2c - a - b) > 1$ and where

$$c_1 := a(2c - a) + b(2c - b) - 2c + 1 - \frac{ab}{d}(4c - a - b - 1)$$

and $c_2 := \frac{4}{d}(a + b + 1) - 8$.

The result (1.1) is an extension of the following classical Gauss’ second summation theorem (see, e.g., [1, p. 11], [6, p. 69] and [7, p. 350]):

$${}_2F_1 \left[\begin{matrix} a, b; \\ \frac{1}{2}(a + b + 1); \end{matrix} \frac{1}{2} \right] = \frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}a + \frac{1}{2}\right)\Gamma\left(\frac{1}{2}b + \frac{1}{2}\right)} \tag{1.3}$$

and the result (1.2) is an extension of the following classical Watson’s summation theorem (see, e.g., [1, p. 16] and [7, p. 351]):

$${}_3F_2 \left[\begin{matrix} a, b, c; \\ \frac{1}{2}(a + b + 1), 2c; \end{matrix} 1 \right] = \frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(c + \frac{1}{2}\right)\Gamma\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)\Gamma\left(c - \frac{1}{2}a - \frac{1}{2}b + \frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}a + \frac{1}{2}\right)\Gamma\left(\frac{1}{2}b + \frac{1}{2}\right)\Gamma\left(c - \frac{1}{2}a + \frac{1}{2}\right)\Gamma\left(c - \frac{1}{2}b + \frac{1}{2}\right)} \tag{1.4}$$

$(\Re(2c - a - b) > -1)$.

Here and in the following, each of denominator parameters in the ${}_pF_q$ is assumed to be nonzero.

Kim et al. [4] established the summation formula (1.2) by using some results contiguous to Gauss’ second summation theorem obtained by Lavoie et al. [5]. Here, we provide a new proof of the extended Watson’s summation theorem (1.2). We also consider some interesting special cases of (1.2). For an interesting proof of the classical Watson’s summation theorem (1.4), the reader may be referred to [3].

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/321643396>

NOTE ON THE EXTENDED WATSON'S SUMMATION THEOREM FOR THE SERIES

Article in *Far East Journal of Mathematical Sciences* · December 2017

DOI: 10.17654/MS102112895

CITATION

1

READS

434

3 authors:



Junesang Choi

Dongguk University-Gyeongju Campus

407 PUBLICATIONS 5,420 CITATIONS

[SEE PROFILE](#)



Vivek Rohira

1 PUBLICATION 1 CITATION

[SEE PROFILE](#)



Arjun Kumar Rathie

Vedant College of Engineering & Technology (Rajasthan Technical University), Bundi ...

267 PUBLICATIONS 1,377 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Hypergeometric Functions and Their Applications [View project](#)



Series involving the zeta functions [View project](#)

2. Derivation of (1.2)

Let \mathcal{L} be the left side of (1.2). Expressing ${}_4F_3$ as the series, we have

$$\mathcal{L} = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k (d+1)_k}{\left(\frac{1}{2}(a+b+3)\right)_k (d)_k 2^k k!} \left\{ \frac{2^k (c)_k}{(2c)_k} \right\}, \quad (2.1)$$

where $(\lambda)_n$ is the Pochhammer symbol defined (for $\lambda \in \mathbb{C}$) by (see [7, p. 2 and pp. 4-6]):

$$\begin{aligned} (\lambda)_n &:= \begin{cases} 1 & (n = 0) \\ \lambda(\lambda+1)\cdots(\lambda+n-1) & (n \in \mathbb{N}) \end{cases} \\ &= \frac{\Gamma(\lambda+n)}{\Gamma(\lambda)} \quad (\lambda \in \mathbb{C} \setminus \mathbb{Z}_0^-), \end{aligned} \quad (2.2)$$

where Γ is the familiar Gamma function. Here and in what follows, let \mathbb{C} , \mathbb{N} and \mathbb{Z}_0^- be the sets of complex numbers, positive integers and non-positive integers, respectively, and let $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. Applying a known identity (see, e.g., [6, p. 49])

$${}_2F_1 \left[\begin{matrix} -\frac{1}{2}k, -\frac{1}{2}k + \frac{1}{2}; \\ c + \frac{1}{2}; \end{matrix} ; 1 \right] = \frac{2^k (c)_k}{(2c)_k} \quad (\Re(c) > 0; k \in \mathbb{N}_0)$$

to (2.1), we obtain

$$\mathcal{L} = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k (d+1)_k}{\left(\frac{1}{2}(a+b+3)\right)_k (d)_k 2^k k!} {}_2F_1 \left[\begin{matrix} -\frac{1}{2}k, -\frac{1}{2}k + \frac{1}{2}; \\ c + \frac{1}{2}; \end{matrix} ; 1 \right]. \quad (2.3)$$

It easily follows from (2.2) that

$$(\lambda)_{2n} = 2^{2n} \left(\frac{1}{2}\lambda\right)_n \left(\frac{1}{2}\lambda + \frac{1}{2}\right)_n \quad (n \in \mathbb{N}_0). \quad (2.4)$$

Using (2.4), we get

$$\left(-\frac{1}{2}k\right)_m \left(-\frac{1}{2}k + \frac{1}{2}\right)_m = 2^{-2m}(-k)_{2m} = \frac{2^{-2m}k!}{(k-2m)!} \quad (k, m \in \mathbb{N}_0). \quad (2.5)$$

Expressing ${}_2F_1$ in (2.3) as the series with the aid of (2.5), we obtain

$$\mathcal{L} = \sum_{k=0}^{\infty} \sum_{m=0}^{\lfloor \frac{k}{2} \rfloor} \frac{(a)_k (b)_k (d+1)_k 2^{-2m}}{\left(\frac{1}{2}(a+b+3)\right)_k 2^k (d)_k \left(c + \frac{1}{2}\right)_m m! (k-2m)!}. \quad (2.6)$$

Applying the following formal manipulation of double series (see, e.g., [2] and [6, p. 57, Lemma 11]):

$$\sum_{k=0}^{\infty} \sum_{m=0}^{\lfloor \frac{k}{2} \rfloor} A(m, k) = \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} A(m, k+2m)$$

to (2.6) and using the identity

$$(a)_{k+2m} = (a)_{2m} (a+2m)_k$$

in the resulting double infinite series, we have

$$\begin{aligned} \mathcal{L} &= \sum_{m=0}^{\infty} \frac{(a)_{2m} (b)_{2m} (1+d)_{2m}}{\left(\frac{k}{2}(a+b+3)\right)_{2m} 2^{4m} (d)_{2m} \left(c + \frac{1}{2}\right)_m m!} \\ &\times \sum_{k=0}^{\infty} \frac{(a+2m)_k (b+2m)_k (d+1+2m)_k}{\left(\frac{1}{2}(a+b+3)+2m\right)_k (d+2m)_k 2^k k!}. \end{aligned} \quad (2.7)$$

Expressing the inner series in (2.7) as a ${}_3F_2$, we get

$$\mathcal{L} = \sum_{m=0}^{\infty} \frac{(a)_{2m} (b)_{2m} (1+d)_{2m}}{\left(\frac{k}{2}(a+b+3)\right)_{2m} 2^{4m} (d)_{2m} \left(c + \frac{1}{2}\right)_m m!}$$

$$\times {}_3F_2 \left[\begin{matrix} a + 2m, b + 2m, 1 + d + 2m; \\ \frac{1}{2}(a + b + 3) + 2m, d + 2m; \end{matrix} \frac{1}{2} \right]. \quad (2.8)$$

Now, we can evaluate the ${}_3F_2$ series in (2.8) with the help of (1.1), after simplification, using Gauss' second summation theorem (1.3), we arrive at the right side of (1.2).

3. Special Cases of (1.2)

Here, we consider some interesting and potentially useful special cases of (1.2).

(i) First, setting $b = -2n$ and replacing a by $a + 2n$ and second, letting $b = -2n - 1$ and substituting $a + 2n + 1$ for a ($n \in \mathbb{N}_0$) in (1.2), we see that, in each case, one of the two terms on the right side of the resulting identities (1.2) will vanish. We thus have

$$\begin{aligned} & {}_4F_3 \left[\begin{matrix} -2n, a + 2n, c, d + 1; \\ \frac{1}{2}(a + 3), 2c, d; \end{matrix} 1 \right] \\ &= \frac{\alpha(a + 1)}{(2c - a - 1)(a + 4n + 1)(a + 4n - 1)} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2} - c + \frac{1}{2}a\right)_n}{\left(\frac{1}{2}a + \frac{1}{2}\right)_n \left(c + \frac{1}{2}\right)_n}, \end{aligned} \quad (3.1)$$

where $n \in \mathbb{N}_0$ and

$$\alpha := (a + 2n)(2c - a - 2n) - 4n(c + n) - 2c + 1 + \frac{2n(a + 2n)}{d}(4c - a - 1);$$

$$\begin{aligned} & {}_4F_3 \left[\begin{matrix} -2n - 1, a + 2n + 1, c, d + 1; \\ \frac{1}{2}(a + 3), 2c, d; \end{matrix} 1 \right] \quad (n \in \mathbb{N}_0) \\ &= \left(1 - \frac{a + 1}{2d}\right) \frac{2(a + 1)}{(a + 4n + 1)(a + 4n + 3)} \frac{\left(\frac{3}{2}\right)_n \left(\frac{3}{2} - c + \frac{1}{2}a\right)_n}{\left(\frac{1}{2}a + \frac{1}{2}\right)_n \left(c + \frac{1}{2}\right)_n}. \end{aligned} \quad (3.2)$$

(ii) Also, setting $d = \frac{1}{2}(a + 1)$ and $d = 2c - 1$ in (3.1), respectively, we obtain the following summation formulas:

$${}_3F_2 \left[\begin{matrix} -2n, a + 2n, c; \\ \frac{1}{2}(a + 1), 2c; \end{matrix} \middle| 1 \right] = \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2} - c + \frac{1}{2}a\right)_n}{\left(\frac{1}{2}a + \frac{1}{2}\right)_n \left(c + \frac{1}{2}\right)_n} \quad (3.3)$$

and

$${}_3F_2 \left[\begin{matrix} -2n, a + 2n, c; \\ \frac{1}{2}(a + 3), 2c - 1; \end{matrix} \middle| 1 \right] = \beta \frac{\left(\frac{1}{2}\right)_n \left(\frac{3}{2} - c + \frac{1}{2}a\right)_n}{\left(\frac{1}{2}a + \frac{1}{2}\right)_n \left(c - \frac{1}{2}\right)_n}, \quad (3.4)$$

where $n \in \mathbb{N}_0$ and

$$\beta := \frac{(a + 1)(a - 1)}{(a + 4n + 1)(a + 4n - 1)}.$$

(iii) Further, setting $d = \frac{1}{2}(a + 1)$ and $d = 2c - 1$ in (3.2), respectively, we get

$${}_3F_2 \left[\begin{matrix} -2n - 1, a + 2n + 1, c; \\ \frac{1}{2}(a + 1), 2c; \end{matrix} \middle| 1 \right] = 0 \quad (3.5)$$

and

$${}_3F_2 \left[\begin{matrix} -2n - 1, a + 2n + 1, c; \\ \frac{1}{2}(a + 3), 2c - 1; \end{matrix} \middle| 1 \right] = \gamma \frac{\left(\frac{3}{2}\right)_n \left(\frac{1}{2}a - c + \frac{3}{2}\right)_n}{\left(\frac{1}{2}a + \frac{1}{2}\right)_n \left(c + \frac{1}{2}\right)_n}, \quad (3.6)$$

where $n \in \mathbb{N}_0$ and

$$\gamma := \frac{(a + 1)(4c - a - 3)}{(a + 4n + 1)(a + 4n + 3)(2c - 1)}.$$

We note that the identities (3.3)-(3.6) have been presented by Lavoie et al. [5] who employed other methods different from those used here. We also remark that the method of proof used here is potentially useful in getting some other summation formulas for the ${}_pF_q$.

Acknowledgment

The authors would like to express their deep-felt thanks for the reviewer's helpful comments.

References

- [1] W. N. Bailey, Generalized Hypergeometric Series, Cambridge University Press, Cambridge, 1935; Reprinted by Stechert Hafner, New York, 1964.
- [2] J. Choi, Notes on formal manipulations of double series, Commun. Korean Math. Soc. 18(4) (2003), 781-789.
- [3] J. Choi and P. Agarwal, Note on the classical Watson's theorem for the series ${}_3F_2$, Honam Math. J. 35(4) (2013), 701-706.
- [4] Y. S. Kim, M. A. Rakha and A. K. Rathie, Extensions of certain classical summation theorems for the series ${}_3F_1$, ${}_3F_2$ and ${}_4F_3$ with applications in Ramanujan's summations, Int. J. Math. Math. Sci. 2010 (2010), Article ID 309503, 26 pp.
- [5] J. L. Lavoie, F. Grondin and A. K. Rathie, Generalizations of Watson's theorem on the sum of a ${}_3F_2$, Indian J. Math. 34(2) (1992), 23-32.
- [6] E. D. Rainville, Special Functions, Macmillan Company, New York, 1960; Reprinted by Chelsea Publishing Company, Bronx, New York, 1971.
- [7] H. M. Srivastava and J. Choi, Zeta and q -Zeta Functions and Associated Series and Integrals, Elsevier Science Publishers, Amsterdam, London and New York, 2012.